

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Microbiota and volatilome profile of fresh and chill-stored deepwater rose shrimp (*Parapenaeus longirostris*)**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1727747> since 2020-02-14T23:34:17Z

*Published version:*

DOI:10.1016/j.foodres.2020.109057

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**1 Microbiota and volatilome profile of fresh and chill-stored deepwater**  
**2 rose shrimp (*Parapenaeus longirostris*)**

**3**  
**4 Foteini F. Parlapani<sup>a,\*</sup>, Ilario Ferrocino<sup>b</sup>, Sofia Michailidou<sup>c</sup>, Anagnostis**  
**5 Argiriou<sup>c</sup>, Serkos A. Haroutounian<sup>d</sup>, Lambros Kokokiris<sup>e</sup>, Kalliopi Rantsiou<sup>b</sup>,**  
**6 Ioannis S. Boziaris<sup>a</sup>**

**7**  
**8 <sup>a</sup> Lab. Marketing and Technology of Aquatic Products and Foods, Dept. of**  
**9 Ichthyology and Aquatic Environment, School of Agricultural Sciences, University of**  
**10 Thessaly, Fitokou street, 38446, N. Ionia, Volos, Greece**

**11 <sup>b</sup> DISAFA, Microbiology and Food Technology Sector, University of Turin, Largo**  
**12 Paolo Braccini 2, 10095 Grugliasco, Italy**

**13 <sup>c</sup> Institute of Applied Biosciences, Centre for Research and Technology Hellas**  
**14 (CERTH), 57001, Thessaloniki, Greece**

**15 <sup>d</sup> Department of Animal Science and Aquaculture, Agricultural University of Athens,**  
**16 Iera Odos 75, 118 55, Athens, Greece**

**17 <sup>e</sup> Department of Nutritional Sciences and Dietetics, International Hellenic University**  
**18 (former Alexander Technological and Educational Institute of Thessaloniki), 57400,**  
**19 Sindos, Thessaloniki, Greece**

**20**  
**21**  
**22 \* Corresponding author. Mailing address: School of Agricultural Sciences, Dept of**  
**23 Ichthyology and Aquatic Environment, Fitokou street, 38446, N. Ionia, Volos,**  
**24 Greece. Phone: +30-24210-93137. Fax: +30-24210-93157. e-mail: fwparlap@uth.gr**

## ABSTRACT

Bacterial communities and Volatile Organic Compounds (VOCs) profile of deepwater rose shrimp (*Parapenaeus longirostris*) stored at 0°C (ice) and 4°C were investigated using 16S amplicon based sequencing and Solid phase micro-extraction (SPME) - Gas chromatography/mass spectrometry (GC/MS), respectively. The shelf-life of shrimps determined by sensory assessment was 5 and 2 days at 0°C and 4°C, respectively. Based on 16S analysis (culture-independend), the initial microbiota of shrimps mainly consists of *Photobacterium*, *Candidatus Hepatoplasma*, *Psychrobacter*, *Acinetobacter* and *Delftia*. *Psychrobacter* and *Carnobacterium* dominated during storage at both temperatures. *Psychrobacter* was the most dominant taxon at the end of shelf-life of chill-stored shrimps. A minor microbial population composed by *Brevundimonas*, *Stenotrophomonas*, *Staphylococcus*, *Legionella*, *Acinetobacter*, *Bacillus*, *Escherichia-Shigella*, *Enterococcus*, *Enterobacter*, *Klebsiella* was also detected. Those taxa may be originated from the environment due to an inadequate hygienic practice during fishing, handling and icing. VOCs such as ethanol, 3-methyl-1-butanol, 2-ethyl-1-hexanol, 3-hydroxy-2-butanone, indole etc., were found to be associated with shrimps at 4°C, while acetone and dimethyl sulfide with shrimps in ice. Some VOCs, from microbial or chemical origin, increased in shrimps either at 0°C (i.e. 1-octen-3-ol, *trans*-2-octenal) or at 4°C (i.e. 3-methyl-1-butanol, indole), while 2-methylbutanal and 3-methylbutanal increased in both temperatures. A positive correlation between *Psychrobacter* with 2-ethyl-1-hexanol and *Carnobacterium* with 3-methyl-1-butanol was also observed. Concluding, we suggest the reinforcement of Good Hygiene Practices on fishing boats during fishing/handling, the rapid onboard icing and keeping shrimps iced avoiding even small increase of storage temperature that affects quality parameters (e.g. microbial

population level, synthesis of microbiota, VOCs profile) in order to provide a product of the highest quality and safety in the market.

**Keywords:** crustaceans, shrimps, spoilage, microbiota, 16S Next Generation Sequencing, Volatile Organic Compounds

## 1. Introduction

Crustaceans are traded and consumed in million tons around the world. The deepwater rose shrimp (*Parapenaeus longirostris*, Lucas 1846) is an important commercial species of the coasts of Italy, Spain, France, Algeria, Tunisia, Greece and Turkey, with Italy (4631 tons) and Spain (4237 tons) to be the countries with the largest catches (FAO, 2019). *P. longirostris* is one of the most landed and highly consumed shrimp species in Greece, where it is consumed as unprocessed. However, rose shrimp is extremely sensitive to deterioration even when it is stored in ice (shelf-life 6 days, Mendes, Gonçalves, Pestana & Pestana, 2005).

Despite the spoilage of shrimp begins with the appearance of melanosis (Ashie, Smith & Simpson, 1996; Lopez-Caballero, Martinez-Alvarez, Gomez-Guillen & Montero, 2006), microorganisms grow quickly and deteriorate its quality. Fresh shrimp spoils due to the rich content of protein, non-protein nitrogen compounds and other nutrients that can allow microbial growth (Don, Xavier, Devi, Nayak & Kannuchamy, 2018; Licciardello, Kharchoufi, Muratore & Restuccia, 2018). Microbial spoilage of shrimp begins immediately after catch and the microbial growth can be controlled by rapid icing of shrimps (Heinsz, Harrison & Leiting, 1988). A consortium of bacteria, the so-called Specific Spoilage Organisms (SSOs), is a small part of the initial total microbiota which grows faster over the other members of the

total microbiota and produces metabolites responsible for the development of off-odors and off-flavors during storage resulting the organoleptic rejection of the product (Gram & Huss, 1996). The domination of particular microorganisms or SSOs depends on various factors such as fish species, geographical origin, synthesis of the initial fish microbiota, microbial interactions, contamination during fishing, handling, storage and distribution in combination with the applied storage conditions (Bozaris & Parlapani, 2016; Parlapani et al., 2018b). The type of the microorganisms that prevail during storage determines the type of fish spoilage e.g. the production of Volatile Organic Compounds (VOCs) and the development of specific off-odors and off-flavors (Bozaris & Parlapani, 2016).

The use of 16S Next Generation Sequencing is able to reveal cultivable and non-cultivable bacterial genera directly from the sample which they may be involved in spoilage of shrimps (Yang, Xie & Qian, 2017) and other seafood (Chaillou et al. 2015; Kuuliala et al. 2018; Parlapani et al. 2018a; 2018b; 2019; Zotta, Parente, Ianniello, De Filippis & Ricciardi, 2019). In addition, microorganisms may be associated with hygiene conditions during fishing, icing, landing and distribution or potential pathogens coming from various sources of contamination can be also highlighted (Parlapani et al. 2018b). The derived information can be used to implement preventive measures for the improvement of microbial quality and safety of fish during handling and storage.

Despite the increasing consumption of deepwater rose shrimp, there is no study, to our knowledge, about bacterial communities and volatile compounds production during chill storage. The aim of this study was: (i) to determine microbiota through 16S amplicon sequencing, (ii) to investigate the VOCs profile using SPME-GC/MS and (iii) to correlate the volatilome with microbiome and sensory score for

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

101 chill-stored shrimps, in order to reveal the initial microbiota of shrimps, the  
102 microbiota changes during storage and the produced VOCs at the time point the  
103 particular microorganisms dominate the shrimp causing spoilage.

## 104 105 **2. Materials and Methods**

### 106 *2.1 Shrimps provision, handling and storage*

107 Two different batches of the deepwater rose shrimp (*P. longirostris*), that were  
108 caught by trawl fishery from Strymonikos Gulf (mean weight $\pm$ sd: 12.3 $\pm$ 3.0g, 72m  
109 depth, F/V Athanassios) in April (10-04-2017, 17-04-2017) were packaged in  
110 insulated boxes with melted ice (0°C) and transferred to the Laboratory of Marketing  
111 and Technology of Aquatic Products and Foods (Department of Ichthyology and  
112 Aquatic Environment, University of Thessaly, Volos, Greece) within four hours from  
113 capture. The insulated boxes containing 1kg of shrimps each, were stored in  
114 incubators operating at two different temperature conditions; 0°C (shrimps with  
115 melted ice replaced every day) and 4°C.

### 116 117 *2.2 Evaluation of shrimp's shelf-life*

118 Sensory evaluation was performed by 5 trained panelists, who evaluated the  
119 appearance of the shell (bright colours, slight blackening on the head, blackening on  
120 head and body), and flesh odor (fresh, marine, musty, ammoniacal, sour, putrid),  
121 according to FAO/WHO (1999). The sensory attributes rating were scored using a 5  
122 to 1 scale with 5 (excellent), 4 (good), 3 (acceptable), 2 (unacceptable) and 1 (spoiled)  
123 scores. A score of 3 was considered as the score for minimum acceptability and the  
124 time point that average score was below 3 (which means that at least one out of the  
125 five panelists scored with 2) was considered as the rejection time point.

126

### 127 *2.3 Microbiological analysis*

128 Ten grams (10 g) flesh from 10 shrimp individuals were aseptically transferred  
129 into stomacher bags with 90 mL MRD (Maximum Recovery Diluent, 0.1% w/v  
130 peptone, 0.85% w/v NaCl) and homogenized for 2 min using a Stomacher (Bug  
131 Mixer, Interscience, London, UK). Using the spread plate technique, 0.1 mL from 10-  
132 fold serial dilutions were used in order to enumerate the total microbial population as  
133 aerobic plate counts (APC) on TSA (Tryptone Soy Agar) and *Pseudomonas* on  
134 ceftrimide-fucidin-cephaloridine agar (CFC) after incubation at 25°C for 48 h. Using  
135 the pour plating method with overlay technique, 1 mL of serial dilutions in MRD was  
136 used for the enumeration of counts on Iron Agar (IA) by counting only the black  
137 colonies (H<sub>2</sub>S producing bacteria) after incubation at 25°C for 72 h, Lactic Acid  
138 Bacteria (LAB) on De Man, Rogosa, Sharpe agar (MRS) after incubation at 25°C for  
139 72 h and counts on Violet Red Bile Glucose agar (VRBGA) by counting only the red  
140 to dark purple colonies surrounded by a reddish zone (Enterobacteriaceae), after  
141 incubation at 37°C for 24 h. The results were expressed as mean log cfu g<sup>-1</sup> ± standard  
142 deviation (sd) of 4 replicates (two replicates per batch).

143

### 144 *2.4 16S rRNA meta-barcoding analysis*

#### 145 *DNA extraction, quality evaluation and sequencing*

146 Twenty five (25) grams of flesh per batch (N=20 individuals) were used to  
147 collect bacterial pellets for 16S rRNA amplicon sequencing, as it was described in  
148 Parlapani et al. (2018b). For each sample, 200 µl of each diluted pellet were used for  
149 bacterial DNA extraction with ZR Soil Microbe DNA MicroPrep (ZYMO  
150 RESEARCH; Irvine, CA, USA). DNA concentration was measured on a Qubit 2.0

151 Fluorimeter using the Qubit® dsDNA BR assay kit (Invitrogen, Carlsbad, CA, USA).

152 The total DNA (DNA from two batches) was pooled before the analysis.

153 Metagenomic analysis was performed by amplifying the V3–V4 region of the  
154 16S rRNA gene using the Illumina’s 16S Metagenomic Sequencing Library  
155 Preparation (15044223 B) protocol. For the amplification of the V3–V4 region, gene-  
156 specific primers were selected based on the Klindworth et al. (2013), by adding  
157 Illumina overhang adapter nucleotide sequences at the 5’ end. PCR reactions were  
158 performed in a 36-well rotor carousel on a Rotor-Gene Q Thermocycler (Qiagen,  
159 Hilden, Germany) according to Parlapani et al. (2018b).

160 PCR products were purified to remove unincorporated primers and primer-  
161 dimer species using NucleoMag® NGS Bead Suspension (Macherey-Nagel, Düren,  
162 Germany). A second PCR was performed to attach dual indices and Illumina  
163 sequencing adapters in all PCR fragments following the instruction of the Illumina’s  
164 16S Metagenomic Sequencing Library Preparation. All libraries were quantified with  
165 fluorometric quantification using Qubit® dsDNA BR assay kit and their molarity was  
166 calculated in relation to the size of DNA amplicons after indexing. Quantitative PCR  
167 (qPCR) was conducted on a Rotor-Gene Q thermocycler (Qiagen, Hilden, Germany)  
168 with the KAPA Library Quantification kit for Illumina sequencing platforms (KAPA  
169 BIOSYSTEMS, Woburn, MA, U.S.A.).

170 Sequencing was performed on a MiSeq platform according to the  
171 manufacturer’s instructions, using the MiSeq® reagent kit v3 (2 x 300 cycles)  
172 (Illumina, San Diego, California).

173

174 *Bioinformatics and Data Analysis*



Trim Galore wrapper (Krueger, 2015) was used for quality trimming of raw reads (.fastq files). Sequences were clustered into operational taxonomic units (OTUs) using the Quantitative Insights into Microbial Ecology (QIIME) pipeline (Caporaso et al., 2010), with 99% sequence similarity to SILVA 128 release for QIIME database. OTU picking was executed using the default clustering algorithm UCLUST (Edgar, 2010). Further processing of OTU tables was conducted in R version 3.4.2 (R Development Core Team, 2010). OTU counts and taxonomic assignments were merged to a phyloseq object with phyloseq R package (McMurdie & Holmes, 2013). Sequences have been submitted through the NCBI sequencing submission portal and deposited under the SRA: SUB6377492, with accession number SAMN12878864.

#### *2.5 VOCs determination by Headspace SPME-GC/MS analysis*

Ten (10) g of flesh from approximately ten shrimps were homogenized. Five grams of the homogenized sample in duplicate per batch were transferred into a 20 mL glass vial (two replicates per batch, 2x2=4 replicates in total). The SPME-GC/MS analysis was performed according to Parlapani et al. (2015a). Compounds were identified by comparing: (i) the linear retention indices (LRI) based on an homologous series of even numbered n-alkanes (C8–C24, Niles, Illinois, USA) with those of standard compounds and literature data, and (ii) the MS data with those of reference compounds and that of MS data obtained from NIST library (NIST/EPA/NIH Mass Spectral Library with Search Program, data version NIST 05, software version 2.0d). The Amdis software (version 2.62, <http://chemdata.nist.gov/mass-spc/amdis/>) was used for the deconvolution of mass spectra and the identification of target components. The amount of VOCs was

expressed in arbitrary units of the peak area of the deconvoluted component multiplied by  $10^{-6}$ .

## 2.6 Statistical analysis

For the 16S data plots were visualized by combining functions provided by the ggplot2 R package (Wickham, 2009).

Alpha diversity indices were calculated using the Chao1 and ACE indices (Chao & Chiu, 2016) were assessed to measure species richness and the Shannon, Simpson, Inverse Simpson and Fisher indices (Lande, 1996) were used to assess the proportional abundances and frequencies of the identified species by the *vegan* package of R.

Abundance estimations for each species were normalized to 100% within each sample, thus, percentages do not reflect the true biomass fraction of each sample. Venn diagrams were constructed using the InteractiVenn online tool (Heberle, Meirelles, da Silva, Telles & Minghim, 2015) to depict pairwise comparisons of shared, common and/or unique OTUs. Spearman's non-parametric correlations were used through the R package *psych* to study the relationships between microbial taxa abundance and volatilome profile and visualized in R using the *corrplot* package of R.

Differences of means in microbial populations were subjected to Analysis of Variance (ANOVA) followed by Tukey's significant difference test, using STATISTICA 6.0. A probability level of  $p \leq 0.05$  was considered statistically significant.

## 3. Results

### 3.1 Rejection time of chill-stored shrimps

Sensory attributes of shrimps were evaluated as ‘excellent’ at the beginning of the experiments (T0, Table 1). Their shell presented bright colors. Flesh odor was fresh and marine. At the time of minimum acceptability (grade 3), the shrimp odor was weak "fishy odour", even slight ammonia was produced, and the color was natural light pink with grey-greenish, and the shell lacked luster sheen. After this point, the shrimps exhibited unpleasant odor, thus they were graded as unfit and rejected (Table 1). Therefore, the shelf-life determined by sensory assessment was estimated to be 5 days at 0°C (ice) and 2 days at 4°C.

### 3.2 Microbial population changes

At the beginning of the shrimp’s shelf-life (T0), the APC was found to be  $3.12 \pm 0.58$  log cfu/g, while the populations of *Pseudomonas*, H<sub>2</sub>S producing bacteria and LAB were found to be  $2.69 \pm 0.61$ ,  $2.04 \pm 0.08$ ,  $3.00 \pm 0.70$  log cfu/g, respectively (Table 2). During storage, the bacteria were grown faster at 4°C reaching 4 to 5 logs for all the population monitored after one day of storage ( $p \leq 0.05$ , T1, Table 2) and reached up to 6.50 logs after two days of storage (T2). In respect of ice-stored shrimps, APC was found to be  $6.59 \pm 0.36$  log cfu/g after five days of storage (T5), while the populations of *Pseudomonas*, H<sub>2</sub>S producing bacteria and LAB reached  $6.11 \pm 0.53$ ,  $4.96 \pm 0.63$  and  $5.23 \pm 0.47$  log cfu/g respectively (Table 2). Enterobacteriaceae remained below the detection limit of 1 log cfu/g throughout the experiment.

### 3.3 NGS analysis

#### 3.3.1 Illumina MiSeq data analysis

After sequencing 500,206 reads were obtained and after the quality filtering 465,369 sequences were assigned to 4,026 unique OTUs Table S1. In particular, 90.6%, 90.4%, 89.4%, 88.5% and 85.1% of reads were assigned to the Phylum, Class, Order, Family and Genus level, respectively. The most diverse community was observed for T2-0°C storage conditions in which the highest number of unique OTUs was observed among the studied samples (Table S1). That was also verified by the Chao1 index (Table S2).

### 3.3.2. Microbiota composition

A simple microbiota was observed in all the samples analysed with the predominance of *Photobacterium* (18.0%) as the main microbes in fresh shrimps (T0), while *Psychrobacter* and *Carnobacterium* dominated together in shrimps during storage at both temperatures (Table 3). *Carnobacterium* was recorded at abundances of 62.4 % at 0° and 20.5% at 4°C at the middle stages of storage, while the *Psychrobacter* at abundances of 34.0 at 0°C and 52.5% at 4°C (Table 3). *Psychrobacter* became the most dominant microorganism at the end of shelf-life of shrimps (57.7% and 48.0% in samples at 0°C and 4°C, respectively), while *Carnobacterium* was also found at 29.8% and 41.2% at 0°C and 4°C, respectively (Table 3). Other genera like *Candidatus Hepatoplasma*, *Pseudomonas*, *Acinetobacter*, *Delftia*, *Brevundimonas*, *Stenotrophomonas*, *Bacillus*, *Vibrio*, *Vagococcus* were also found but at lower abundances (up to 11.3%) in fresh or chill-stored samples (Table 3), while other 167 genera including *Staphylococcus*, *Legionella*, *Bacillus*, *Escherichia-Shigella*, *Enterococcus*, *Enterobacter*, *Klebsiella* etc. were found at abundances of less than 1% in all cases (Table S3).

### 3.4 VOCs produced in chill-stored shrimps

A total of 41 VOCs were identified (11 alcohols, 12 aldehydes, 11 ketones, 3 esters and other 4 compounds: dimethyl sulfide, methylene chloride, carbon disulfide and indole). By comparing the presence of the VOCs between T0 and samples stored in ice or in refrigerate condition it was possible to observe that samples stored at 4°C were characterized by the presence of ethanol, 3-methyl-1-butanol, 2-ethyl-1-hexanol, *trans*-2-nonenal, *trans*-2,*cis*-6-nonadienal, 3-hydroxy-2-butanone, 3-octen-2-one and indole, while acetone and dimethyl sulfide were mainly associated with ice storage (Fig 1a). The compounds 1-heptanol, 1-nonanol, octanal, 3-pentanone and methylene chloride were associated with the fresh shrimps (T0, Fig 2a). Some compounds were also increased in shrimps stored at 0°C (i.e. 1-octen-3-ol and *trans*-2-octenal) or at 4°C (i.e. 3-methyl-1-butanol, *trans*-2,*cis*-6-nonadienal and indole), while the compound 2-methylbutanal and 3-methylbutanal were increased at both 0 and 4°C (Fig 1b).

### 3.5 Correlation between Microbiome and Volatilome in rose shrimps

Spearman's correlation between the microbial OTUs and VOCs clearly indicated the correlation of the most abundant bacterial species in rose shrimps with the concentration of metabolites produced during their chill storage ( $P < 0.05$ , Fig 2). *Psychrobacter* and *Carnobacterium* which found to be the most abundant genera during storage at both temperatures, they were positively correlated with some VOCs. In particular, *Psychrobacter* was positively correlated with 2-ethyl-1-hexanol, 2-nonanone and 2-octanone, and *Carnobacterium* with 3-methyl-1-butanol and 2-pentanone. Positive correlations were also found between *Candidatus Hepatoplasma* and 1-heptanol, 1-nonanol, octanal, 3-pentanone, *Photobacterium* and 2-butanone,

298 *Vagococcus* and 3-methyl-1-butanol, 2-methylbutanal, 3-methylbutanal, 2-octanone,  
299 2-nonanone, *Vibrio* and 2-butanone, ethyl acetate (Fig 2). On the other hand,  
300 *Psychrobacter* was negatively correlated with dimethyl sulphide, while *Delftia* and  
301 *Lactobacillus* with 3-methyl-1-butanol (Fig 2).

302

#### 303 **4. Discussion**

304 The 16S sequencing coupled with VOCs profile analysis were used to  
305 investigate microbial spoilage of the deepwater rose shrimp caught from Greek  
306 seawaters. The correlation of the microbiome with volatilome and their link with  
307 sensory score allow us to understand how the quality of shrimp is being deteriorated  
308 during storage and how it is eventually rejected.

309 Fresh shrimp remains one of the most delicious and delicatessen food in  
310 restaurants, catering and all-inclusive hotels along Greek coasts, inland and islands.  
311 However, the shelf-life of the deepwater rose shrimp is short as it was determined in  
312 this study (5 days in ice and 2 days at 4°C). *Pseudomonas*, H<sub>2</sub>S producing bacteria  
313 and LAB are usually responsible for the quality loss of chill-stored shrimp such as the  
314 tropical brackish water shrimp *Penaeus notialis* (Dabadé et al., 2015). In the present  
315 study, these organisms reached the levels of 5-6 logs at the end of fish shelf-life, with  
316 *Pseudomonas* to dominate the shrimp using the classical microbiological approach.  
317 However, *Pseudomonas* was found at very low abundances using the 16S NGS  
318 analysis (culture independed approach). This indicates that organisms other than  
319 *Pseudomonas* might grew on CFC culture medium. The microbiota grown on the  
320 selective culture media has to be investigated using molecular methodologies in future  
321 studies.

Microorganisms such as *Acinetobacter*, *Bacillus* and *Delftia*, which were found during the first days of storage, are usually associated with the handling conditions and contamination occurs from the human skin, equipment, soil, washing water or ice, etc. (de Amorim & Nascimento, 2017; Heinsz et al., 1988; Parlapani et al., 2018). Other bacteria coming from human and environmental sources like *Enterobacter*, *Escherichia*, *Shigella*, *Klebsiella*, *Legionella*, *Enterococcus* and *Staphylococcus* are also involved in the inadequate attention to hygiene during fishing, handling and icing affecting fish quality and safety. The fact that these genera were found at abundances less than 1% herein does not minimize the possibility of pathogens presence. All of them contain human pathogens, foodborne or not, might be found in raw shrimps. In that case, a likely consumption of undercooked shrimps could cause foodborne infections causing socioeconomic hardship in the Greek market, like legal penalties, shutdown of business, and tarnished reputation. On the other hand, the detection of them using 16S NGS does not indicate potential risk. NGS is not able to differentiate living and dead cells, thus overestimating the living taxa in microbial communities (Emerson et al., 2017). To minimize contamination in order to maximize the quality and safety of fishery products, the findings derived from this and previous studies (Parlapani et al. 2018a; 2018b; 2019) could be used to develop approaches for rapid analysis like the High Resolution Melting analysis (HRM) or the multiplex PCR using specific genes as targets for the identification of pathogenic species/strains contained into the genera identified using 16S metabarcoding. In addition, we could strengthen extension activities (education, communication and outreach) for individuals, communities and industries involved in the primary production in Greece.

346 The 16S data also revealed herein the presence of *Candidatus Hepatoplasma*,  
 347 *Photobacterium* and *Psychrobacter* at the beginning of deepwater rose shrimp shelf-  
 348 life. From those, *Candidatus Hepatoplasma* has been described as symbionts of the  
 349 terrestrial isopods *Porcellio scaber* and *Oniscus asellus* (Wang, Stingl, Anton-  
 350 Erxleben, Zimmer & Brune, 2004; Wang, Brune & Zimmer, 2007), while the other  
 351 two bacteria are usually found in fresh fish (Gram & Huss 1996). *Photobacterium* has  
 352 been also reported as spoiler of fish under particular storage conditions.  
 353 *Photobacterium* (*P. phosphoreum* or *Photobacterium* sp.) usually predominate in fish  
 354 such as the cod, the halibut and the cold-smoked salmon under elevated CO<sub>2</sub>  
 355 concentrations (Modified Atmosphere Packaging – MAP) or vacuum (Gram &  
 356 Dalgaard, 2002; Hovda, Sivertsvik, Lunestad, Lorentzen & Rosnes, 2007; Kuuliala et  
 357 al. 2018; Olofsson, Ahrne & Molin, 2007; Stohr, Joffraud, Cardinal & Leroi, 2001).  
 358 The facultative fermentative metabolism of *Photobacterium* and the competitive  
 359 interactive environment defined by the presence of other bacteria in fish flesh stored  
 360 under aerobic conditions (Gram & Dalgaard, 2002) could may explain the absence of  
 361 its growth increase during the storage of the deepwater rose shrimp. *Psychrobacter*  
 362 and *Carnobacterium* were observed to co-dominate during storage in this study, at the  
 363 first two days at 4°C, and the last two days at 0°C. This means that *Psychrobacter* and  
 364 *Carnobacterium* bacteria dominate the shrimps stored at the higher temperature (4°C),  
 365 in a shorted time than the ice-stored shrimps (0°C). *Psychrobacter* usually dominate  
 366 the chill-stored crustaceans like the brown shrimp (Broekaert, Heyndrickx, Herman,  
 367 Devlieghere & Vlaemynck, 2013a; Broekaert, Nosedá, Heyndrickx, Vlaemynck &  
 368 Devlieghere, 2013b), the Pacific white shrimp (Yang et al., 2017), the Norway lobster  
 369 (Bekaert, Devriese, Maes & Robbens, 2015), but also the fish like the gilt-head sea  
 370 bream (Parlapani et al., 2018a), the thawed hake, the thawed plaice fillets (Zotta et



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

371 al., 2019) or even the cephalopods like the thawed cuttlefish (Parlapani et al., 2018b).  
372 This bacterium might causes the spoilage of the brown shrimp through the break  
373 down of lipids and the hydrolysis of proteins and amino acids (Broekaert et al.,  
374 2013b). On the other hand, *Carnobacterium* was found to dominate the peeled brown  
375 shrimp stored under MAP at 4°C (Calliauw et al., 2016a), or it was detected in  
376 shrimps stored in MAP (Dalgaard et al., 2003) or in fish species stored in air  
377 (Parlapani et al., 2018a; Zotta et al., 2019).

378         The compound 3-methyl-1-butanol has been associated with the metabolic  
379 activity of *Carnobacterium* in inoculated cooked and peeled tropical shrimp *Penaeus*  
380 *vannamei* stored under MAP at 8°C (Jaffres et al., 2011). In our study, this compound  
381 was positively correlated to the presence of the *Carnobacterium* and it was increased  
382 in shrimps stored at 4°C. *C. divergens* and *C. maltaromaticum* are known to cause  
383 sensory spoilage of cooked and peeled shrimps stored under MAP at 5°C, producing  
384 ammonia, tyramine, alcohols (including 3-methyl-1-butanol), aldehydes and ketones  
385 (including 2-pentanone, Laursen, Leisner, & Dalgaard, 2006). Other compounds such  
386 as the ethanol, the dimethyl sulfide and the ethyl acetate were being increased in  
387 cooked peeled gray/brown shrimp *Crangon crangon* stored under MAP (Nosedá et  
388 al., 2012). The ethyl acetate, which showed a positive correlation with *Vibrio* herein,  
389 has been also reported to the brown shrimp (Kuuliala et al., 2018). *Vagococcus*  
390 (particularly *V. salmoninarum*) was found to produce 2-nonanone in inoculated brown  
391 shrimp stored under MAP at 4°C (Calliauw, Horemans, Broekaert, Michiels &  
392 Heyndrickx, 2016b). From the aforementioned compounds, only the 3-methyl-1-  
393 butanol increased in this study. Additionally, the 1-octen-3-ol and the *trans*-2-octenal  
394 were increased in the deepwater rose shrimps stored at 0°C, the *trans*-2,*cis*-6-  
395 nonadienal and indole in those stored at 4°C, while the 2-methylbutanal and the 3-

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

396 methylbutanal they were increased in the shrimps stored at both temperatures. The 1-  
397 octen-3-ol, *trans*-2-octenal and *trans*-2,*cis*-6-nonadienal are associated with chemical  
398 oxidations in fish (Duflos, Coin, Cornu, Antinelli & Malle, 2006; Iglesias & Medina,  
399 2008), while the 2-methylbutanal and 3-methylbutanal are products of microbial  
400 activity (Parlapani, Mallouchos, Haroutounian & Boziaris, 2017). The compounds 3-  
401 methyl-1-butanol, 1-octen-3-ol, *trans*-2-octenal, 2-methylbutanal, 3-methylbutanal  
402 and indole they also exhibited an increasing trend in chill-stored blue crabs and  
403 proposed as potential spoilage markers of that product (Parlapani et al., 2019). The  
404 indole has been associated with the spoilage of *P. longirostris* (Mendes et al., 2005;  
405 Mendes, Huidobro & Caballero, 2002) and it has been found as a product of various  
406 bacteria like *Vibrio* spp. and *Morganella morganii* isolated from the wild shrimps  
407 *Litopenaeus setiferus* and *Litopenaeus brasiliensis* (Benner, Staruszkiewicz & Otwell,  
408 2004). Herein, no correlation between indole (which found at very low relative  
409 concentrations at 4°C) and bacteria was observed. This may occurs since indole is not  
410 produced in significant quantities by bacteria in shrimps stored at low temperatures  
411 (Mendes et al., 2002). Other compounds such as hexanol, heptanol, hexanal, heptanal,  
412 octanal and nonanal found in rose shrimp are known as aroma compounds and are  
413 mainly coming from enzymatic reactions, oxidation or autoxidation of lipids  
414 (Alasalvar, Taylor & Shahidi, 2005; Duflos et al., 2006).

415 Overall, the inadequate hygiene during fishing/handling can cause microbial  
416 contamination from human, foodstuffs and environmental sources. In this way, the  
417 initial total microbial population increases, the quality of fish product is lost rapidly  
418 and the product might not be safe to consume. For these reasons, fishermen have to  
419 reinforce the Good Hygiene Practices regarding the potential sources of microbial  
420 contamination during handling which usually reduce quality and shorten shelf-life of

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

421 fish. Moreover, the shrimps have to be stored and distributed in ice (rapid onboard  
422 icing and keeping fish iced until marketing or consumption). To reinforce the  
423 competitiveness of fishery products at local, national or international commercial  
424 level, the research outputs should be communicated through websites, workshops,  
425 public dialogues, newsletters, and one-on-one consultations in order to create  
426 awareness among academia, fishery sectors and society.

## 427 428 **Conclusions**

429 The 16S meta-barcoding revealed microorganisms associated with safety or spoilage  
430 of the deepwater rose shrimp from Greek seawaters. The simulated storage conditions  
431 showed how the higher temperature causes the fast domination of the shrimps by the  
432 spoilers *Psychrobacter* and *Carnobacterium* resulting to their rapid quality  
433 deterioration (shelf-life: 2 days at 4°C vs 5 days in ice). In addition, compounds  
434 associated with microbial activity e.g. 3-methyl-1-butanol, 2-ethyl-1-hexanol and 2-  
435 nonanone presented a positive correlation with the dominant organisms e.g.  
436 *Psychrobacter* or *Carnobacterium* in rose shrimp. From them, 3-methyl-1-butanol  
437 increased during storage. Other compounds associated with chemical reactions e.g. 1-  
438 octen-3-ol, *trans*-2-octenal and *trans*-2,*cis*-6-nonadienal were also increased. This  
439 indicates that not only microbial action, but also chemical mechanisms might be  
440 responsible for the quality loss of chill-stored rose shrimp. Based on our findings, we  
441 could suggest preventive measures to improve the quality and safety of rose shrimp,  
442 particularly fishermen have to reinforce the Good Hygiene Practices during fishing  
443 and handling, to accomplish rapidly onboard icing and keep shrimps iced avoiding  
444 even small changes in storage temperature that affect the microbial population level,  
445 synthesis of microbiota, VOCs profile and eventually the shelf-life of the product. The

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

446 results of this study will be used to prepare a guide for the proper handling and  
447 storage conditions of the fishing products produced by the small-scale fishery sector  
448 in Greece in order to provide a product of the highest quality and safety in the market.

## 450 **Acknowledgments**

451 This research has been co-financed by the European Union (European Social Fund —  
452 ESF) and the Greek government through the Operational Program “IKY Scholarship  
453 Programs for Strengthening Post Doctoral Research” in the frame of the Human  
454 Resources Development Program, Education and Lifelong Learning.

## 456 **5. References**

457 Alasalvar, C., Taylor, K.D.A. & Shahidi, F. (2005). Comparison of volatiles of  
458 cultured and wild sea bream (*Sparus aurata*) during storage in ice by dynamic  
459 headspace analysis/Gas Chromatography–Mass Spectrometry. *Journal of Agricultural*  
460 *and Food Chemistry*, 53, 2616-2622.

461  
462 Ashie, I. N. A., Smith, J. P. & Simpson, B. K. (1996). Spoilage and shelf-life extension of  
463 fresh fish and shellfish. *Critical Reviews in Food Science and Nutrition*, 36, 87-121.

464  
465 Arlinghaus, R., Tillner, R., & Bork, M. (2015). Explaining participation rates in  
466 recreational fishing across industrialised countries. *Fisheries Management and*  
467 *Ecology*, 22, 45–55.

468  
469 Bekaert, K., Devriese, L., Maes, S., & Robbens, J. (2015). Characterization of the  
470 dominant bacterial communities during storage of Norway lobster and Norway lobster

471 tails (*Nephrops norvegicus*) based on 16S rDNA analysis by PCR-DGGE. *Food*  
 472 *Microbiology*, 46, 132-138.  
 473  
 474 Benner, R. A., Jr., Staruszkiewicz, W. F., & Otwell, W. S. (2004). Putrescine,  
 475 cadaverine, and indole production by bacteria isolated from wild and aquacultured  
 476 penaeid shrimp stored at 0, 12, 24, and 36 C. *Journal of Food Protection*, 67(1), 124–  
 477 133.  
 478  
 479 Bozianis, I. S., & Parlapani, F. F. (2016). Specific Spoilage Organisms (SSO) in Fish.  
 480 In A. Bevilacqua, M. R. Corbo, M. Sinigaglia & R. Sykes (Eds.), *Microbiological*  
 481 *Quality of Food: Foodborne Spoilers* (pp 60-98). Woodhead Publishing.  
 482  
 483 Broekaert, K., Heyndrickx, M., Herman, L., Devlieghere, F., & Vlaemynck, G.  
 484 (2013a). Molecular identification of the microbiota of peeled and unpeeled brown  
 485 shrimp (*Crangon crangon*) during storage on ice and at 7.5 °C. *Food Microbiology*,  
 486 36, 123–134.  
 487  
 488 Broekaert, K., Nosedá, B., Heyndrickx, M., Vlaemynck, G., & Devlieghere, F.  
 489 (2013b). Volatile compounds associated with *Psychrobacter* spp.  
 490 and *Pseudoalteromonas* spp., the dominant microbiota of brown shrimp (*Crangon*  
 491 *crangon*) during aerobic storage. *International Journal of Food Microbiology*, 166,  
 492 487-493.  
 493  
 494 Calliauw, F., De Mulder, T., Broekaert, K., Vlaemynck, G., Michiels, C., &  
 495 Heyndrickx, M. (2016a). Assessment throughout a whole fishing year of the dominant

496 microbiota of peeled brown shrimp (*Crangon crangon*) stored for 7 days under  
 497 modified atmosphere packaging at 4 °C without preservatives. *Food Microbiology*,  
 498 54, 60-71.  
 499  
 500 Calliauw, F., Horemans, B., Broekaert, K., Michiels, C., & Heyndrickx, M. (2016b)  
 501 Spoilage potential of *Vagococcus salmoninarum* in preservative-free, MAP-stored  
 502 brown shrimp and differentiation from *Brochothrix thermosphacta* on streptomycin  
 503 thallous acetate actidione agar. *Journal of Applied Microbiology*, 120(5), 1302-1312.  
 504  
 505 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello,  
 506 E.K., et al. (2010). QIIME allows analysis of high-throughput community sequencing  
 507 data. *Nature Methods*, 7, 335–336. 10.1038/nmeth.f.303.  
 508  
 509 Chaillou, S., Chaulot-Talmon, A., Caekebeke, H., Cardinal, M., Christieans, S.,  
 510 Denis, C., et al., (2014). Origin and ecological selection of core and food-specific  
 511 bacterial communities associated with meat and seafood spoilage. *The ISME Journal*,  
 512 1–14.  
 513  
 514 Chao A., & Chiu, C. (2016). Species richness: estimation and comparison Wiley  
 515 StatsRef: Statistics reference online (pp. 1–26). New York, NY: John Wiley & Sons,  
 516 Ltd.  
 517  
 518 Dabadé, D. S., den Besten, H. M., Azokpota, P., Nout, M. R., Hounhouigan, D. J., &  
 519 Zwietering, M. H. (2015). Spoilage evaluation, shelf-life prediction, and potential

520 spoilage organisms of tropical brackish water shrimp (*Penaeus notialis*) at different  
 521 storage temperatures. *Food Microbiology*, 48, 8–16.  
 522  
 523 Dalgaard, P., Vancanneyt, M., Euras Vilalta, N., Swings, J., Fruekilde, P., & Leisner,  
 524 J. (2003). Identification of lactic acid bacteria from spoilage associations of cooked  
 525 and brined shrimps stored under modified atmosphere between 0 °C and 25 °C.  
 526 *Journal of Applied Microbiology*, 94(1), 80–9.  
 527  
 528 Dalgaard, P. (1995). Qualitative and quantitative characterization of spoilage bacteria  
 529 from packed fish. *International Journal of Food Microbiology*, 26, 319-333.  
 530  
 531 De Amorim, A. M. B., & Nascimento, J. S. (2017) Acinetobacter: an underrated  
 532 foodborne pathogen? *Journal of Infection in Developing Countries*, 11(2), 111-114.  
 533  
 534 Don, S., Xavier, K.A.M., Devi, S.T., Nayak, B.B., & Kannuchamy, N. (2018).  
 535 Identification of potential spoilage bacteria in farmed shrimp (*Litopenaeus vannamei*):  
 536 Application of Relative Rate of Spoilage models in shelf life prediction. *LWT - Food*  
 537 *Science and Technology*, 97, 295–301.  
 538  
 539 Duflos, G., Coin, V.M., Cornu, M., Antinelli, J.F., & Malle, P. (2006). Determination  
 540 of volatile compounds to characterize fish spoilage using headspace/mass  
 541 spectrometry and solid-phase microextraction/ gas chromatography/mass  
 542 spectrometry. *Journal of the Science of Food and Agriculture*, 86, 600–611.  
 543

544 Edgar, R.C. (2010). Search and clustering orders of magnitude faster than BLAST.  
 545 *Bioinformatics*, 26(19), 2460–2461.  
 546  
 547 Emerson, J.B., Adams, R.I., Román, C.M., Brooks, B., Coil, D.A., Dahlhausen, K.,  
 548 Ganz, H.H., Hartmann, E.M., Hsu, T., Justice, N.B., & Paulino-Lima, I.G. (2017).  
 549 Schrödinger's microbes: tools for distinguishing the living from the dead in microbial  
 550 ecosystems. *Microbiome*, 5, 86.  
 551  
 552 FAO (2016). The State of World Fisheries and Aquaculture 2016. Contributing to  
 553 food security and nutrition for all. Rome. 200 pp.  
 554  
 555 FAO (2018). The State of World Fisheries and Aquaculture 2018 - Meeting the  
 556 sustainable development goals. Rome. Licence: CC BY-NC-SA 3.0 IGO.  
 557  
 558 FAO. Fisheries and Aquaculture Department. (2019).  
 559 [http://www.fao.org/fishery/species/2598/en/Accessed 23 September 2019](http://www.fao.org/fishery/species/2598/en/Accessed%2023%20September%202019).  
 560  
 561 FAO/WHO (1999). Guidelines for the sensory evaluation of fish and shellfish in  
 562 laboratories. CAC-GL 31-1999.  
 563  
 564 Gram, L., & Huss, H. H. (1996). Microbiological spoilage of fish and fish products.  
 565 *International Journal of Food Microbiology*, 33, 121-137.  
 566  
 567 Gram, L. & Dalgaard, P. (2002). Fish spoilage bacteria – problems and solutions.  
 568 *Current Opinion in Biotechnology*, 13, 262-266.



569

570 Heberle, H.; Meirelles, G. V.; da Silva, F. R.; Telles, G. P.; Minghim, R. (2015).

571 *InteractiVenn: a web-based tool for the analysis of sets through Venn diagrams. BMC*

572 *Bioinformatics*, 16, 169.

573

574 Heinsz, L. J., Harrison, M. A., & Leiting, V. A. (1988). Microflora of brown shrimp

575 (*Penaeus aztecus*) from Georgia coastal waters. *Food Microbiology*, 5, 141-145.

576

577 Hovda, M. B., Sivertsvik, M., Lunestad, B. T., Lorentzen, G., & Rosnes, J. T. (2007).

578 Characterisation of the dominant bacterial population in modified atmosphere

579 packaged farmed halibut (*Hippoglossus hippoglossus*) based on 16S rDNA-DGGE.

580 *Food Microbiology*, 24, 362–37.

581

582 Jaffrès, E., Lalanne, V., Macé, S., Cornet J., Cardinal M., Sérot T., Dousset,

583 X., Joffraud, J.-J. (2011). Sensory characteristics of spoilage and volatile compounds

584 associated with bacteria isolated from cooked and peeled tropical shrimps using

585 SPME–GC–MS analysis. *International Journal of Food Microbiology*, 147(3), 195-

586 202.

587

588 Iglesias, J. & Medina, I. (2008). Solid-phase microextraction method for the

589 determination of volatile compounds associated to oxidation of fish muscle. *Journal*

590 *of Chromatography A*, 1192, 9e16.

591

592 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., &

593 Glöckner, F.O. (2013). Evaluation of general 16S ribosomal RNA gene PCR primers

594 for classical and next-generation sequencing-based diversity studies. *Nucleic Acids*  
595 *Res.*, 41, e1.

596

597 Kuuliala, L., Al Hage, Y., Ioannidis, A.-G., Sader, M., Kerckhof, F.-M., Vanderroost,  
598 M., Boon, N., De Baets, B., De Meulenaer, B., Ragaert, P., & Devlieghere, F. (2018).  
599 Microbiological, chemical and sensory spoilage analysis of raw Atlantic cod (*Gadus*  
600 *morhua*) stored under modified atmospheres. *Food Microbiology*, 70, 232-244.

601

602 Lande, R. (1996). Statistics and partitioning of species diversity, and similarity among  
603 multiple communities. *Oikos*, 76, 5-13.

604

605 Laursen, B.G., Leisner, J.J., & Dalgaard, P. (2006). Carnobacterium Species: Effect of  
606 Metabolic Activity and Interaction with *Brochothrix thermosphacta* on Sensory  
607 Characteristics of Modified Atmosphere Packed Shrimp. *Journal of Agricultural*  
608 *and Food Chemistry*, 54, 3604-3611.

609

610 Licciardello, F., Kharchoufi, S., Muratore, G., & Restuccia, C. (2018). Effect of  
611 edible coating combined with pomegranate peel extract on the quality maintenance of  
612 white shrimps (*Parapenaeus longirostris*) during refrigerated storage. *Food*  
613 *Packaging and Shelf Life*, 17, 114–119.

614

615 Lopez-Caballero, M. E., Martinez-Alvarez, O., Gomez-Guillen, M. C. & Montero, P  
616 (2006). Effect of natural compounds alternative to commercial antimelanosis on  
617 polyphenol oxidase activity and microbial growth in cultured prawns (*Marsupenaeus*  
618 *tiger*) during chill storage. *European Food Research Technology*, 223, 7-15.

619

620 McMurdie, P. J., & Holmes, S., (2012). phyloseq: A Bioconductor Package for  
621 Handling and Analysis of High-Throughput Phylogenetic Sequence Data.  
622 *Pacific Symposium on Biocomputing*, 17, 235–246.

623

624 Mendes, R., Gonçalves, A., Pestana, J., & Pestana, C. (2005). Indole production and  
625 deepwater pink shrimp (*Parapenaeus longirostris*) decomposition. *European Food*  
626 *Research and Technology*, 221(3-4), 320-328.

627

628 Mendes, R., Huidobro, A., & Caballero, E. L. (2002). Indole levels in deepwater pink  
629 shrimp (*Parapenaeus longirostris*) from the Portuguese coast. Effects of temperature  
630 abuse. *European Food Research and Technology*, 214(2), 125-130.

631

632 Nosedá, B., Goethals, J., De Smedt, L., Dewulf J., Samapundo S., Van Langenhove,  
633 H., & Devlieghere, F. (2012). Effect of O<sub>2</sub>/CO<sub>2</sub> enriched atmospheres on  
634 microbiological growth and volatile metabolite production in packaged cooked peeled  
635 gray shrimp (*Crangon crangon*). *International Journal of Food Microbiology*, 160(1),  
636 65-75.

637

638 Olofsson, T. C., Ahrne, S., & Molin, G. (2007). The bacterial flora of vacuum-packed  
639 cold-smoked salmon stored at 7°C, identified by direct 16S rRNA gene analysis and  
640 pure culture technique. *Journal of Applied Microbiology*, 103, 1364-5072.

641

642 Parlapani, F. F., Anagnostopoulos, D. A., Koromilas, S., Kios, K., Michailidou, S.,  
643 Pasentis, K., Psomopoulos, F., Argiriou, A., Haroutounian, S.A., & Boziaris I. S.

644 (2019). Bacterial communities and potential spoilage markers of whole Blue Crab  
 645 (*Callinectes sapidus*) stored under commercial simulated conditions. *Food*  
 646 *Microbiology*, 82, 325-333.  
 647  
 648 Parlapani, F.F., Malouchos, A., Haroutounian, S.A., & Boziaris, I.S. (2017).  
 649 Microbial and non-microbial origin volatile organic compounds produced on model  
 650 fish substrate inoculated or not with gilt-head sea bream spoilage bacteria. *LWT -*  
 651 *Food Science and Technology*, 78, 54-62.  
 652  
 653 Parlapani, F. F., Michailidou, S., Anagnostopoulos, D. A., Sakellariou, A. K.,  
 654 Pasentsis, K., Psomopoulos, F., Argiriou, A., Haroutounian, S. A., & Boziaris, I. S.  
 655 (2018a). Microbial spoilage investigation of thawed common cuttlefish (*Sepia*  
 656 *officinalis*) stored at 2°C using next generation sequencing and volatilome analysis.  
 657 *Food Microbiology*, 76, 518–525.  
 658  
 659 Parlapani, F.F., Michailidou, S., Pasentsis, K., Argiriou, A., Krey, G., & Boziaris, I.S.  
 660 (2018b). A meta-barcoding approach to assess and compare the storage temperature  
 661 dependent bacterial diversity of gilt-head sea bream (*Sparus aurata*) originating from  
 662 fish farms from two geographically distinct areas of Greece. *International Journal of*  
 663 *Food Microbiology*, 278, 36–43.  
 664  
 665 R Development Core Team (2010). R: A language and environment for statistical  
 666 computing, R Foundation for Statistical Computing, Vienna, Austria.  
 667

668 Stohr, V., Joffraud, J.J., Cardinal, M., & Leroi, F. (2001). Spoilage potential and  
 669 sensory profile associated with bacteria isolated from cold-smoked salmon. *Food*  
 670 *Research International*, 34(9), 797-806.  
 671  
 672 Wang, Y., Brune, A., & Zimmer, M. (2007). Bacterial symbionts in the  
 673 hepatopancreas of isopods: diversity and environmental transmission. *FEMS*  
 674 *Microbiology Ecology*, 61(1), 141-152.  
 675  
 676 Wang, Y., Stingl, U., Anton-Erxleben, F., Zimmer, M., & Brune, A. (2004).  
 677 'Candidatus Hepatincola porcellionum' gen. nov., sp. nov., a new, stalk-forming  
 678 lineage of Rickettsiales colonizing the midgut glands of a terrestrial isopod. *Archives*  
 679 *of Microbiology*, 181(4), 299-304.  
 680  
 681 Wickham, H. (2009). ggplot2: elegant graphics for data analysis. Springer; New York.  
 682  
 683 Yang, S.-P., Xie, J., & Qian, Y.-F. (2017). Determination of spoilage microbiota of  
 684 Pacific white shrimp during ambient and cold storage using Next-Generation  
 685 Sequencing and culture-dependent method. *Journal of Food Science*, 82, 1178-1183.  
 686  
 687 Zotta, T., Parente, E., Ianniello, R.G., De Filippis, F., & Ricciardi, A. (2019).  
 688 Dynamics of bacterial communities and interaction networks in thawed fish fillets  
 689 during chilled storage in air. *International Journal of Food Microbiology*, 293, 102-  
 690 113.  
 691  
 692

693

694

695

696

697

698

699

700

701

702

703

704 **Table 1.** Sensory score of deepwater rose shrimp stored at 0°C (ice) and 4°C.

	0°C (ice)				4°C	
Sensory score	T0	T2	T4	T5	T1	T2
<i>Shell appearance</i>	Excellent	Good	Acceptable	Unacceptable	Acceptable	Unacceptable
<i>Flesh odor</i>	Excellent	Excellent	Good	Unacceptable	Good	Unacceptable

705

706

707 **Table 2.** Population changes (mean log cfu g<sup>-1</sup> ± sd) of four replicates during storage

708 (in days, T) of shrimps stored at 0°C (ice) and 4°C.

	0°C (ice)				4°C	
	T0	T2	T4	T5	T1	T2
<b>APC</b>	3.12±0.58 <sup>a</sup>	4.32±0.38 <sup>b</sup>	5.92±0.38 <sup>c</sup>	6.59±0.36 <sup>d</sup>	4.85±0.62 <sup>b</sup>	6.44±0.46 <sup>c</sup>
<i>Pseudomonas</i>	2.69±0.61 <sup>a</sup>	3.44±0.43 <sup>b</sup>	5.39±0.46 <sup>c</sup>	6.11±0.53 <sup>d</sup>	4.46±0.51 <sup>e</sup>	5.97±0.47 <sup>c</sup>

<b>H<sub>2</sub>S bacteria</b>	2.04±0.08 <sup>a</sup>	2.82±0.57 <sup>b</sup>	4.14±0.73 <sup>c</sup>	4.96±0.63 <sup>d</sup>	4.15±0.49 <sup>c</sup>	5.14±0.10 <sup>d</sup>
--------------------------------	------------------------	------------------------	------------------------	------------------------	------------------------	------------------------

<b>LAB</b>	3.00±0.70 <sup>a</sup>	3.69±0.85 <sup>a</sup>	4.43±0.33 <sup>b</sup>	5.23±0.47 <sup>c</sup>	4.29±0.61 <sup>b</sup>	5.08±0.48 <sup>c</sup>
------------	------------------------	------------------------	------------------------	------------------------	------------------------	------------------------

Mean values with different letters are significantly different (p<0.05). Mean population values along different time (T0-T5) or temperature values (0 and 4°C), are compared using ANOVA.

**Table 3.** The Relative abundance (%) of the dominant genera assigned to 16S rRNA sequences, in the deepwater rose shrimp stored in 0 and 4°C.

OTUs (%)	0°C (ice)				4°C	
	T0	T2	T4	T5	T1	T2
<i>Psychrobacter</i>	8.10	20.9	34.0	57.7	52.5	48.0
<i>Carnobacterium</i>	1.60	8.70	62.4	29.8	20.5	41.2
<i>Pseudomonas</i>	0.30	7.30	0.60	0.70	11.3	0.60
<i>Photobacterium</i>	18.0	0.10	0.00	0.00	0.00	0.20
<i>Acinetobacter</i>	5.20	0.90	0.20	5.30	0.10	0.10
<i>Delftia</i>	4.20	6.70	0.10	0.10	0.10	0.00
<i>Candidatus Hepatoplasma</i>	10.0	0.00	0.00	0.00	0.00	0.00
<i>Vagococcus</i>	0.00	0.00	0.60	2.00	1.90	5.30
<i>Stenotrophomonas</i>	1.10	7.00	0.00	0.20	0.20	0.30
<i>Brevundimonas</i>	6.50	1.50	0.10	0.30	0.10	0.10
<i>Bacillus</i>	0.30	6.90	0.00	0.30	0.00	0.10
<i>Vibrio</i>	5.40	0.00	0.00	0.00	0.00	0.20
<i>Exiguobacterium</i>	0.10	0.00	0.00	0.00	4.40	0.00
<i>Lactobacillus</i>	4.30	0.10	0.00	0.00	0.10	0.00
<i>Paucimonas</i>	1.50	2.20	0.00	0.10	0.00	0.00
<i>Planococcus</i>	0.10	3.20	0.00	0.30	0.20	0.10

1		<i>Rhizobium</i>	1.00	2.30	0.00	0.10	0.10	0.00
2		<i>Aerococcus</i>	0.10	0.20	0.00	0.10	2.60	0.30
3	720							
4								
5	721							
6								
7	722							
8								
9								
10	723							
11								
12								
13	724							
14								
15	725							
16								
17	726							
18								
19								
20	727							
21								
22								
23								
24								
25								
26								
27								
28								
29								
30								
31								
32								
33								
34								
35								
36								
37								
38								
39								
40								
41								
42								
43								
44								
45								
46								
47								
48								
49								
50								
51								
52								
53								
54								
55								
56								
57								
58								
59								
60								
61								
62								
63								
64								
65								



## Legends to Figures

**Figure 1. Panel A. Abundance of VOCs (relative concentrations, area x 10<sup>-6</sup> under the chromatographic peak) during storage of fresh shrimp (T0, red bars) at Time stored in ice (green bars) or 4°C (blue bars). Panel B. Shift in volatilome profile during storage at different storage conditions (T2, T4 and T5 for ice-stored shrimps and T1\_4°C and T2\_4°C for shrimps stored at 4°C). Boxes represent the interquartile ranges (IQRs) between the first and third quartiles, and the lines inside represent the medians (2nd quartiles). Whiskers denote the lowest and the highest values within IQRs from the first and third quartiles, respectively. Circles represent outliers beyond the whiskers.**

**Figure 2. Spearman's correlation between the bacterial OTUs and VOCs in deepwater rose shrimp. The intensity of the colors indicates the degree of correlation between the bacterial OTUs. The blue color shows a positive correlation, while the red colour shows a negative correlation between OTUs and VOCs.**

1 753  
2  
3 754  
4  
5 755  
6  
7 756  
8  
9 757  
10  
11  
12 758  
13  
14 759  
15  
16 760  
17  
18 761  
19  
20 762  
21  
22 763  
23  
24 764  
25  
26 765  
27  
28 766  
29  
30 767  
31  
32 768  
33  
34 769  
35  
36 770  
37  
38 771  
39  
40 772  
41  
42 773  
43  
44 774  
45  
46 775  
47  
48 776  
49  
50 777  
51  
52 778  
53  
54 779  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**FIG 1-2 (attached file)**

780

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure 1a**  
[Click here to download high resolution image](#)




Figure 1b

[Click here to download high resolution image](#)






Figure 2  
[Click here to download high resolution image](#)











**Foteini F. Parlapani:** Conceptualization, Methodology, Formal analysis, Investigation, Writing-Review and Editing, Project administration

**Ilario Ferrocino:** Methodology, Formal analysis, Investigation, Writing-Review and Editing

**Sofia Michailidou:** Methodology, Formal analysis, Writing-Review and Editing

**Anagnostis Argiriou:** Resources, Writing-Review and Editing

**Serkos A. Haroutounian:** Resources, Writing-Review and Editing

**Lambros Kokokiris:** Resources, Writing-Review and Editing

**Kalliopi Rantsiou:** Resources, Writing-Review and Editing

**Ioannis S. Boziaris:** Resources, Writing-Review and Editing



**UNIVERSITY OF THESSALY**  
**SCHOOL OF AGRICULTURAL SCIENCES**  
**DEPT. ICHTHYOLOGY & AQUATIC ENVIRONMENT**  
**Lab. of Marketing and Technology of Aquatic Products and Foods**  
FITOKO, N. IONIA – 38446 VOLOS, GREECE  
TEL.: +30 24210 93137 / Fax: +30 24210 93157, e-mail: fwparlap@uth.gr

---

Volos, 05-10-2019

Dear Editor,

authors declare no competing interests.

Sincerely Yours,

Dr Foteini F. Parlapani